

Dynamic Field Theory

Part I: continuous spaces and activation fields

Gregor Schöner
gregor.schoener@ini.rub.de

Discrete “neurons”

- or activation variables: how do they arise?
How do they sample sensory/motor spaces...
- no evidence that neural discreteness matters for behavior

Continuity in space

- hypothesis: behavior is embedded in continua
 - the space of possible behaviors, e.g. space of movements, percepts, timing structures
 - neuronal substrate is continuous (maps, broad tuning)
- (\Rightarrow need to understand how categorical behavior may emerge from such continua)

Dynamical Field Theory: space

- in DFT, continuous spaces are dimension over which activation fields are defined
 - homologous to sensory surfaces, e.g., visual or auditory space (retinal, allocentric, ...)
 - homologous to motor surfaces, e.g., saccadic end-points or direction of movement of the end-effector in outer space
 - feature spaces, e.g., localized visual orientations, color, impedance, ...
 - abstract spaces, e.g., ordinal space, along which serial order is represented

example: motion perception

- continuous motion
- apparent motion
- motion pattern

example: selection decisions in motion perception

■ motion pattern

■ why not diagonal motion?



■ or the other diagonal motion?

■ => motion pattern perception involves selection decision


example: selection decisions in motion perception

- can the alternative motion pattern be realized?
 - flat motion quartet
 - tall motion quartet
 - square motion quartet

space of possible percepts and activation field




space of possible actions and activation field



Dynamical Field Theory: space




- fields: continuous activation variables defined over continuous spaces

information, probability, certainty









e.g., retinal space, movement parameters, feature dimensions, viewing parameters, ...

activation fields



representing different percepts




Link between DFT and neurophysiology

- What do neurons represent?


tuning curve

■ example: primary visual cortex (monkey)



tuning curve


■ example: primary motor cortex (monkey)




Link between DFT and neurophysiology

- Example 1: Jancke et al: A17 in the cat, population representation of retinal location



- determine RF profile for each cell
- it's center determines what that neuron codes for
- compute a distribution of population activation by superposing RF profiles weighted with **current** neural firing rate





- The **current** response refers to a stimulus experienced by **all** neurons
- Reference condition: localized points of light



elementary stimuli




- result: population distribution of activation defined over retinal space = representation of visual location



■ => does a decent job estimating retinal position

current stimulus:
square of light

range of retinal field
sampled by neurons



□ 0.4°



- Extrapolate measurement device to new conditions
- e.g., time resolved

two
different
stimulus
locations

time




30 - 40 ms

40 - 50 ms


50 - 60 ms

60 - 70 ms

70 - 80 ms




□ 0.4°



■ or when complex stimuli are presented (here:
two spots of light)

↓ response to composite stimuli increasing distance between the two squares of light
↑



↑ superposition of responses to each
elemental stimulus

□ 0.4°




■ by comparing DPA of composite stimuli to superposition of DPAs of the two elementary stimuli obtain evidence for interaction

■ early excitation


■ late inhibition

interaction




activation level in DPA




at location of left component stimulus




response to composite stimuli



superposition of responses to each elemental stimulus




evidence for inhibitory interaction



time



[ms]

model by dynamic field:




Neurophysiological grounding of DFT

example: movement planning





- tuning of cells in motor and premotor cortex to direction of end-effector movement path



lil023-1



Distribution of Population Activation (DPA)

Distribution of population activation =
 $\sum_{\text{neurons}} \text{tuning curve} * \text{current firing rate}$



■ look at temporal evolution of DPA

■ or DPAs in new conditions, here: DPA reflects prior information



Distributions of Population Activation are abstract

- neurons are not **localized** within DPA!
- cortical neurons really are sensitive to many dimensions
 - motor: arm configuration, force direction
 - visual: many feature dimensions such as spatial frequency, orientation, direction...
- => DPA is a **projection** from that high-dimensional space onto a single dimension