

Basic neurophysics

Gregor Schöner

Institut für Neuroinformatik

SOURCES (except where cited otherwise)


- Peter Dayan and Larry F Abbot: *Theoretical Neuroscience*, MIT Press, Cambridge MA, 2001
 - sections 1.1, 1.2, 1.4, 2.3
- Wulfram Gerstner, Werner M. Kistler, Richard Naud and Liam Paninski: *Neuronal Dynamics: From single neurons to networks and models of cognition*. Cambridge University Press, 2014
 - section 2
 - <http://neurondynamics.epfl.ch/index.html>

the brain

motor
cortex



visual
cortex

frontal
cortex




to motor
output

neurons




$\sim 10^{11}$ with 10000 synapses each

neurons






neurons as input-output units

- inputs from dendrites
- spike formation at soma
- output at axon




two functional components

- membranes: dendrites, soma, axons
- synapses



membrane




source

<http://www.columbia.edu/cu/psychology/courses/1010/mangels/neuro/neurosignaling/neurosignaling.html>

membrane

- membrane=double lipid layer that is an electrical insulator
- neuron is electrically charged: more negative potential inside than outside cell
- based on ions K^+ , Na^+ , and Cl^-




source

<http://www.columbia.edu/cu/psychology/courses/I010/mangels/neuro/neurosignaling/neurosignaling.html>]

membrane


- higher concentration of K^+ inside cell
- lower concentration of Na^+ inside cell
- membrane less permeable to Na^+ than to K^+
 - \Rightarrow Na^+ gradient is steeper than the K^+ gradient
 - \Rightarrow more positive outside cell
 - \Rightarrow negative potential



source

membrane


- gradient comes from ion pumps: protein channels in membrane that transport Na^+ out of cell, K^+ into cell, establishing gradient
- this is where energy is consumed (a lot): ATP used to pump ions



source

membrane


- giant squid axon... used to establish basic biophysics of membrane dynamics
- voltage-clamp



[Source: Neuroscience. 2nd edition. Purves D, Augustine GJ, Fitzpatrick D, et al., editors. Sunderland (MA): Sinauer Associates; 2001.]


synapses

- at a synapse, the membranes of two neurons comes very close
- => this is where transmission across neurons takes place



two types of synapses


- electrical: currents across the membrane directly from one cell to another through “gap junctions”
 - very fast, but not flexible.
 - exists in the peripheral nervous system... but not very common
- chemical: the common one
- that is much more flexible...



[Source: Neuroscience. 2nd edition.
Purves D, Augustine GJ, Fitzpatrick D, et al., editors.
Sunderland (MA): Sinauer Associates; 2001.]

two types of synapses

- chemical: the more common one
- pre-synaptic cell releases neurotransmitter in response to an action potential that arrives through the axon
- post-synaptic potential induced by action of neurotransmitters on receptors




[Source: Neuroscience. 2nd edition.
Purves D, Augustine GJ, Fitzpatrick D, et al., editors.
Sunderland (MA): Sinauer Associates; 2001.]

two types of synapses

■ chemical synapse

- slower transmission... 1 to 2 ms
- but more flexible: tuned by changes in receptors




[Source: Neuroscience. 2nd edition.
Purves D, Augustine G], Fitzpatrick D, et al., editors.
Sunderland (MA): Sinauer Associates; 2001.]

post-synaptic potentials


■ depending on the receptor type, synaptic transmission induces post-synaptic potentials of different forms and sign

■ that travel to the soma, where a spiking decision is made

[Source: Neuroscience. 2nd edition.




spiking mechanism




spiking mechanism

- all or none nature of spikes
- spike generation is coincidence detection
 - overlap of incoming post-synaptic potentials that have propagated to soma within about 10 ms required to sum...
 - typical in cortex: 10 inputs needed, 10000 potential inputs...
- neuron as a “switch”



Hodgkin-Huxley

■ relationship
potential-ionic
concentration



$$\Delta u = u_1 - u_2 = \frac{-kT}{q} \ln \frac{n(u_1)}{n(u_2)}$$

Hodgkin-Huxley

- dynamic model of potential change and three ion currents
- which come from three ion channels
- phenomenological dynamics of the ion ion channels




Fig. 2.2: Schematic diagram for the Hodgkin-Huxley model.

$$C \frac{du}{dt} = - \sum_k I_k(t) + I(t) .$$

$$\sum_k I_k = g_{Na} m^3 h (u - E_{Na}) + g_K n^4 (u - E_K) + g_L (u - E_L) .$$


$$\dot{m} = \alpha_m(u) (1 - m) - \beta_m(u) m$$

$$\dot{n} = \alpha_n(u) (1 - n) - \beta_n(u) n$$


$$\dot{h} = \alpha_h(u) (1 - h) - \beta_h(u) h .$$

Hodgkin-Huxley

■ based on data from squid-axon...




 spikes



Hodgkin Huxley


- the spiking mechanism is an instability => threshold effect




Hodgkin Huxley

- spike rate reflects input current

A




B





Hodgkin Huxley

- time varying inputs make time varying rate



Example: neural circuit

■ stretch reflex



[Source: Neuroscience. 2nd edition.
Purves D, Augustine G], Fitzpatrick D, et al., editors.
Sunderland (MA): Sinauer Associates; 2001.]