Prof. Dr. G. Schöner, Institut für Neuroinformatik

gregor.schoener@rub.de

Autonomous robotics

April 21, 2022

Exercise 1 Dynamical Systems Tutorial

Please upload solutions on the web page before midnight on April 28 (Thursday).

1. Solve the linear differential equation

$$\dot{x} = -\alpha x \tag{1}$$

analytically. If unfamiliar, get help from a text book or online resource.

- 2. Plot the solution as a function of time by setting $\alpha = 1$ /second, using a few discrete times, t = 0, 1, 2, ...10 seconds. Or evaluate the function numerically on the computer in a program of your choice. Make this plot for two values of the initial condition, x(0).
- 3. From the analytical solution, determine the moment in time, τ at which x(t) has fallen to 1/e of its initial value, x(0). Does this time depend on x(0)? Answer the same question for the hyperbolic decay, x(t) = x(0)/(1+t).
- 4. Plot this dynamics Eq (1) and designate the fixed point. Make the same drawing for a dynamics with the same slope and a fixed point at $x_0 > 0$. Write down the equation for that dynamics.
- 5. Make a plot of this nonlinear differential equation

$$\dot{x} = \beta - x^2. \tag{2}$$

Compute its fixed points as a function of β and plot this function. Mark the fixed points in the dynamics as attractors or repellors.